

299 Rio Vista Boulevard, Mermaid Waters 4218 Phone: 07 5575 7922 Fax: 07 5575 8922 Email: admin@goldcoasturologist.com.au

Thursday 24 Oct 2024 Ms Olivia Devet

WHAT TO EXPECT POST YOUR UROLIFT PROCEDURE

This information explains what you may experience post your procedure and how to take care of yourself during this time.

Expect:

- Patients may experience some urinary discomfort during the recovery period
- Blood in the urine which can last up to 2-4 weeks post procedure

• Pain or discomfort when urinating, increased urge and frequency in urination which can last up to 4 weeks

• The catheter is generally removed on the morning after surgery. The expected hospital stay is no more than one night

- If you have had an anaesthetic you will be uninsured to drive for 24hrs
- Return to work is usually 1 week post procedure but this depends on how your recovery is going, type of work and when you are comfortable. Dr Elmes rooms can provide you with a medical certificate if required.

What to do:

- Drink plenty of fluids
- Take Ural for the first 2-3 days if you have penile pain during urination. This helps make your urine more like water and less like lemon juice (maximum sachets a day is 4).

sachets a day is 4)

Movicol is available from your local pharmacy – used to avoid constipation

• Oxytrol OR Vesicare patch - This is used if you have bothersome urgency and frequency, night-time urination incontinence/or penile tip pain. These can be used a maximum of 2 patches at any one time. Side effects: dry mouth,

constipation, dry eyes, forgetfulness and rarely urinary retention. DO NOT use these two medications at the same time.

use these two medications at the same time.

• Endone – prescribed strong pain relieving drug. Side effects: dry mouth, constipation, nausea/vomiting, dizziness, light headed (do not drive if affected)

Avoid:

- Straining/strenuous activity and constipation for 4 weeks
- Driving long distance within first week
- Sexual activity 6 weeks post operatively

Inform Dr Elmes' rooms if:

- Unable to urinate
- Heavy continuous bleeding / multiple clots
- Fevers / unwell